

Isospin and F-spin structure in low-lying levels of $^{48,50}\text{Cr}$ isotopes

F.H. Al-Khudair^{1,2} and G.L. Long^{1,2,3,4}

¹*Department of Physics, Tsinghua University, Beijing 100084, China*

²*Key Laboratory for Quantum Information and Measurements, MOE 100084, China*

³*Center of Nuclear theory, Lanzhou Heavy Ion Accelerator National Laboratory, Lanzhou, 730000, China.*

⁴*Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing, 100080, China*

The low-energy level structure and electromagnetic transitions of $^{48,50}\text{Cr}$ nuclei have been studied by using interacting boson model with isospin (IBM-3). A sequence of isospin excitation bands with isospin ($T = T_z, T_z + 1$ and $T_z + 2$) has been assigned, and compared with available data. According to this study, the 2_3^+ and 2_2^+ states are the lowest mixed symmetry states in $^{48,50}\text{Cr}$ respectively. In particular, the present calculations suggest that a combination of isospin and F-spin excitation can explain the structure in these nuclei. The transition probabilities between the levels are analyzed in terms of isoscalar and isovector decomposition which reveal the detailed nature of the energy levels. The results obtained are found in good agreement with recent experimental data.

I. INTRODUCTION

The interacting boson model (IBM) [1–3] is capable of giving a simple yet realistic description of nuclear collective motions. In its original version (IBM-1), only one kind of boson is considered, corresponding to fully proton neutron symmetric states. The neutron-proton extension of the model (IBM-2), predicts a new class of states [4] having mixed symmetry in the proton and neutron degrees of freedom, and it has been successfully observed in various experiments [5–7].

In lighter nuclei, the valence protons and neutrons fill the same major shell and isospin should be taken into account. In order to include the isospin, IBM has been extended to the interacting boson model with isospin (IBM-3) [8]. In IBM-3 three types of bosons are included: proton-proton (π), neutron-neutron (ν) and proton-neutron (δ) which forms the isospin $T = 1$ multiplet. The ν , δ and π bosons have the isospin projection $M_T = -1, 0, 1$ respectively. The wave functions can be classified by the $U_c(3) \supset SU(2)_T$ group [8], where $SU(2)_T$ is the usual isospin group. Dynamical symmetries of the IBM3 have been studied in Refs. [9,11,13,12,10,14]. The $U_{sd}(6)$ of IBM-1 goes to $U(18)$ group for IBM-3 as its dynamical symmetry group. The natural chains below $U(18)$ start with $U_{sd}(6) \times U_c(3)$, and they must contain $O(3)$ and $SU_T(2)$ as subgroups because the angular momentum and the isospin are good quantum numbers. The chains beginning with $U_{sd}(6) \times U_c(3)$ and satisfying the above requirement are the following [9]

$$\begin{aligned} U(18) &\supset (U_c(3) \supset SU_T(2)) \times (U_{sd}(6) \supset U_d(5) \supset O_d(5) \supset O_d(3)) \\ U(18) &\supset (U_c(3) \supset SU_T(2)) \times (U_{sd}(6) \supset O_{sd}(6) \supset O_d(5) \supset O_d(3)) \\ U(18) &\supset (U_c(3) \supset SU_T(2)) \times (U_{sd}(6) \supset SU_{sd}(3) \supset O_d(3)). \end{aligned} \quad (1)$$

The subgroups $U_d(5)$, $O_{sd}(6)$ and $SU_{sd}(3)$ describe vibrational, γ -unstable and rotational nuclei respectively [15]. The existence of the isospin excitations in light nuclei has received interest in the last few years [16–20]. Such renewed interest was sparked by discovery of states with isospin $T > T_z$ ($T_z = |Z - N|/2$). The Cr isotopes have been the subject of many theoretical and experimental investigations. On the experimental side, large γ -ray detector arrays are now available, and they have increased the detection sensitivity by orders of magnitude. Meanwhile shell model calculations in full fp shell can now be performed. Detailed description of the yrast band structure in the full fp space has been reported in ^{48}Cr [21] and ^{50}Cr [22] respectively. Experimentally, the positive parity yrast band was extended up to the band termination and turned out to be in good agreement with the shell model prediction [23,24]. A detailed analysis of backbending mechanism of ^{48}Cr has been done using projected shell model [25]. In ^{50}Cr the first backbending at $J^\pi = 10^+$ and second backbending at $J^\pi = 18^+$ have been observed experimentally [26]. The high spin states and electromagnetic transitions have been investigated and compared with the shell model results [27]. Recently microscopic three-cluster model calculations were also performed in an attempt to explain the bands structure properties [28]. However a comprehensive analysis of the low-lying levels in these nuclei are still missing. It is the purpose of this paper to study the low-lying levels in the IBM-3, in particular to investigate the isospin and F-spin contents of the levels. This paper is divided as follows. In section II, we briefly discuss the interacting boson model with isospin. In section III, we present the results of our calculation for the energy levels and compared with

available data, and analyze the isospin and F-spin structure of the results. A discussion of electromagnetic transitions follows in section IV. Finally, in section V we summarize our results.

II. THE IBM-3 OPERATORS

The most general IBM-3 Hamiltonian can be written as

$$H = \epsilon_s \hat{n}_s + \epsilon_d \hat{n}_d + H_2, \quad (2)$$

where

$$\begin{aligned} H_2 = & \frac{1}{2} \sum_{L_2 T_2} C_{L_2 T_2} ((d^\dagger d^\dagger)^{L_2 T_2} \cdot (\tilde{d} \tilde{d})^{L_2 T_2}) + \frac{1}{2} \sum_{T_2} B_{0 T_2} ((s^\dagger s^\dagger)^{0 T_2} \cdot (\tilde{s} \tilde{s})^{0 T_2}) \\ & + \sum_{T_2} A_{2 T_2} ((s^\dagger d^\dagger)^{2 T_2} \cdot (\tilde{d} \tilde{s})^{2 T_2}) + \frac{1}{\sqrt{2}} \sum_{T_2} D_{2 T_2} ((s^\dagger d^\dagger)^{2 T_2} \cdot (\tilde{d} \tilde{d})^{2 T_2}) \\ & + \frac{1}{2} \sum_{T_2} G_{0 T_2} ((s^\dagger s^\dagger)^{0 T_2} \cdot (\tilde{d} \tilde{d})^{0 T_2}), \end{aligned} \quad (3)$$

and

$$(b_1^\dagger b_2^\dagger)^{L_2 T_2} \cdot (\tilde{b}_3 \tilde{b}_4)^{L_2 T_2} = (-1)^{(L_2 + T_2)} \sqrt{(2L_2 + 1)(2T_2 + 1)} [(b_1^\dagger b_2^\dagger)^{L_2 T_2} \times (\tilde{b}_3 \tilde{b}_4)^{L_2 T_2}]^{00}, \quad (4)$$

is the dot product in both angular momentum and isospin. The tilted quantity is defined as

$$\tilde{b}_{lm, m_z} = (-1)^{(l+m+1+m_z)} b_{l-m, -m_z}. \quad (5)$$

The symbols T_2 and L_2 represent the two-boson system isospin and angular momentum. The parameters A , B , C , D and G are related to the two-body matrix elements by $A_{T_2} = \langle sd20 | H_2 | sd20 \rangle$, with $T_2 = 0, 1, 2$, $B_{T_2} = \langle s^2 0 T_2 | H_2 | s^2 0 T_2 \rangle$, $G_{T_2} = \langle s^2 0 T_2 | H_2 | d^2 0 T_2 \rangle$, $D_{T_2} = \langle sd 2 T_2 | H_2 | d^2 2 T_2 \rangle$ and $C_{L_2 T_2} = \langle d^2 L_2 T_2 | H_2 | d^2 L_2 T_2 \rangle$, with $T_2 = 0, 2$ and $L_2 = 0, 2, 4$ and by $C_{L_2 1} = \langle d^2 L_2 1 | H_2 | d^2 L_2 1 \rangle$ with $L_2 = 1, 3$. The parameters A_1 , C_{11} and C_{31} are similar to the Majorana interactions in the IBM-2 which will be referred also as Majorana interactions. These interactions are important to shift the states with mixed symmetry with respect to the total symmetric ones. Since only a little experimental information is known about such states in the nuclei under study, we attempt to vary the parameters appearing in these terms to fit the energy of available experimental data on 1^+ and other state which are sensitive to these parameters.

The values of parameters were chosen according to the microscopic studies in Ref [30]. The best fit to the whole spectrum can be found. We have rewritten the Hamiltonian in terms of linear combination of Casimir operators which is convenient to analyze the dynamical symmetry nature. The expressions of the Casimir operators can be found in Ref. [9]. In Casimir operator form, the Hamiltonians are

$$\begin{aligned} H_{48} = & \lambda C_{2U_{sd}(6)} + 1.460T(T+1) + 0.030C_{1U_d(5)} - 0.091C_{2SU_{sd}(3)} \\ & + 0.025C_{2U_d(5)} + 0.173C_{2O_d(5)} + 0.01C_{O_d(3)}, \end{aligned} \quad (6)$$

$$\begin{aligned} H_{50} = & \lambda C_{2U_{sd}(6)} + 1.322T(T+1) + 0.780C_{1U_d(5)} - 0.119C_{2SU_{sd}(3)} \\ & + 0.030C_{2U_d(5)} + 0.091C_{2O_d(5)} + 0.002C_{O_d(3)}, \end{aligned} \quad (7)$$

for $^{48,50}\text{Cr}$ isotopes respectively. The corresponding parameters in the form of eq.(3) are also given in Table I. The λ determines the position of the mixed symmetry states. In ^{50}Cr , there is one 1^+ state at 3.629 MeV, and this requires $\lambda = -0.03$ MeV. We then use the same value for ^{48}Cr isotope because no experimental information is available in this isotope. It can be seen from the Casimir operator form that the Hamiltonian of ^{48}Cr is more rotational than that of ^{50}Cr because the coefficient of $C_{U_d(5)}$ in ^{48}Cr is much less than that in ^{50}Cr . In ^{50}Cr the coefficient of $C_{1U_d(5)}$ is very large, and this indicates that it is more close to the $U(5)$ limit and is in transition from $U(5)$ to $SU(3)$.

III. ISOSPIN AND F- SPIN SYMMETRY STRUCTURE

We assume ^{56}Ni as the closed core and the bosons in the current study corresponding to pairs of hole fermions. The energy levels are shown in Figs. 1 and 2. Good agreement between the calculated and observed spectrum is confirmed up to $J \leq 2N$. The energy levels and wave-functions are obtained using a computer program written by Van Isacker [31]. It is very interesting to see that the sequence of bands with $(T = T_z, T_z + 1 \text{ and } T_z + 2)$ has been produced, and is in a good agreement with available experimental data. The β - band with $T = T_z$ is the first and second excited band in ^{48}Cr and ^{50}Cr respectively. In Fig.1 we also draw the ground state band of ^{48}Ti where the lowest isospin is $T = 2$ which should be close to the isospin analogue state in ^{48}Cr . Here we have assumed that the ground state energy of ^{48}Ti is equal to that of the IBM3 calculated $0_{T=2}^+$ state in ^{48}Cr . This is supported by the following estimate. We estimate the energy of the isospin analogue state in ^{48}Cr by considering the binding energy difference of ^{48}Cr and ^{48}Ti and then subtracting the Coulomb energy difference. This estimation is crude because Coulomb energy depends on the shape of the nucleus sensitively. By using the tables in Ref. [32] and the following Coulomb energy formula

$$E_{Coulomb} = 0.70 \frac{Z^2}{A^{1/3}} (1 - 0.76 Z^{-2/3}), \quad (8)$$

we obtained the energy of the $T = 2$ isospin analogue state in ^{48}Cr to be 8.350 MeV which is close to the energy of the $0_{T=2}^+ = 8.760$ MeV in our IBM-3 calculation.

The energy levels in those figures show that there is good agreement in the ground state and β bands in general. The following points need special attention. In ^{48}Cr , the $3_1^+(T = 0)$ state appears at 6.188 MeV in our calculation, and it is not yet seen in experiment. In ^{50}Cr the IBM-3 predicts the first 3^+ at 3.823 MeV and the second 3^+ at 5.921 MeV. These states are both from the $[N - 1, 1]$ $U(6)$ irreducible representation. Here we have the $U(6)$ labeling as it is a good quantum number approximately. The first and second $J = 4^+$ in $^{48,50}\text{Cr}$ are in good agreement with experimental data, and 4_3^+ state in ^{48}Cr and 4_2^+ state in ^{50}Cr are mixed symmetry states. In both nuclei the first $J = 5^+$ state is a solely mixed symmetry state.

To identify the mixed symmetry states, one can make use of their electromagnetic transition properties: weakly E2 and strong M1 decay to ground state and first 2^+ state respectively [33]. Mixed symmetry $J = 2^+$ state in light nuclei have been identified in $^{54,56}\text{Cr}$, $^{56,58}\text{Fe}$ [34], ^{56}Fe [35,36] and $^{64-68}\text{Ge}$, $^{60-66}\text{Zn}$ [37]. The mixed symmetry structure of wave functions can be seen by calculating the $\langle J | C_{2U(6)} | J \rangle$ value. The calculated 2_3^+ and 2_2^+ in ^{48}Cr and ^{50}Cr exhibit $[N - 1, 1]$ $U(6)$ partition, indicating that those states are the lowest mixed symmetry states in $^{48,50}\text{Cr}$ respectively. For $^{48}\text{Cr}(T_z = 0)$, the lowest mixed symmetry state come from $[N - 2, 2]$ because $[N - 1, 1]$ does not contain $T = 0$, while for ^{50}Cr it comes from $[N - 1, 1]$. From Figs. 1 and 2 one can see this fact and therefore the lowest mixed symmetry state in ^{48}Cr has high energy. The IBM-3 analysis gives a 1_1^+ level at 6.218 MeV with partition $[N - 1, 1]$, and it is higher than the lowest mixed symmetry 2^+ state. No experimental evidence for this level is available. The energy of the first 1^+ in ^{50}Cr equals to 3.613 MeV, and it is quite close to the experimental one. A possible candidate for 1_2^+ mixed symmetry has also been identified at 7.876 MeV, it is close to the observed state $J = (1, 2)$ at 7.646 MeV. The energy of these observed states are well reproduced by the calculation. We have varied each of Majorana parameters around the best-value and keeping all other parameters at their best-fit values, the variations of the energy of these states with the parameters are shown in Figs 3-6. In ^{48}Cr the mixed symmetry component of the lowest mixed symmetry band belongs predominantly to the partition $[N - 2, 2]$ with $T = T_z$. The second mixed symmetry state 2^+ has the partition $[N - 1, 1]$ with $T = T_z + 1 = 1$ with energy 6.110 MeV, it is very close to experimental level at 6.100 MeV ($T = 1$) [41]. Starting with this mixed state, a whole band of mixed states is predicted by this IBM3 calculation. In the same time the lowest mixed symmetry bands in ^{50}Cr have the partition $[N - 1, 1]$ with $T = T_z, T_z + 1$ as shown in Fig. 2. Because IBM-3 has three charge states, it is possible to have $U(6)$ partitions into three rows, namely the $[N_1, N_2, N_3]$ states which are characteristic of IBM-3. We found that such state come high in energy, upwards at about 8.5 MeV, and the lowest example being a scissor mode state in ^{48}Cr at 8.599 MeV which is predominantly the $[2, 1, 1]$ partition with $T = 1$. The present calculation predicts that it decays to 2_3^+ by strongly M1 transition with $B(M1) = 0.24\mu_N^2$ and weakly E2 transition with $B(E2) = 0.0013e^2b^2$. It is very significant if these properties are observed in experiment.

IV. ELECTROMAGNETIC TRANSITION

The E2 transition is described by the following isoscalar and isovector one boson E2 operators $Q = Q^0 + Q^1$ [38]

$$Q^0 = \alpha_0 \sqrt{3} [(s^+ \hat{d})^{20} + (d^+ \hat{s})^{20}] + \beta_0 \sqrt{3} [(d^+ \hat{d})^{20}] \quad (9)$$

$$Q^1 = \alpha_1 \sqrt{2} [(s^+ \hat{d})^{21} + (d^+ \hat{s})^{21}] + \beta_1 \sqrt{2} [(d^+ \hat{d})^{21}]. \quad (10)$$

The M1 transition is also a one boson operator with an isoscalar part and an isovector part $M = M^0 + M^1$ [38],

$$M^0 = g_0 \sqrt{3} (d^+ \hat{d})^{10} = g_0 L / \sqrt{10} \quad (11)$$

$$M^1 = g_1 \sqrt{2} (d^+ \hat{d})^{11}, \quad (12)$$

where g_1 and g_0 are the isovector and isoscalar g -factor respectively and L is angular momentum operator.

Having obtained the wave function for states, we can calculate the electromagnetic transition rate between states using the subroutine of Lac [39]. In order to link it with energy code, we have made some modification. The parameters in the E2 operator are adjusted to fit the experimental data of $B(E2; 2_1^+ \rightarrow 0_1^+)$, where $\alpha_0 = \beta_0 = 0.1, 0.13$ for ^{48}Cr and ^{50}Cr respectively and $\alpha_1 = \beta_1 = 0.1$ for both isotopes. The results are summarized in tables II and III respectively.

The $B(E2)$ values between the yrast states are well reproduced. The calculated $B(E2; 4_1^+ \rightarrow 2_1^+)$ and $B(E2; 6_1^+ \rightarrow 4_1^+)$ are quite close to the experimental values, in the same time the calculated $B(E2; 2_3^+ \rightarrow 0_1^+)$ in ^{50}Cr is larger than the experimental one by about $0.004e^2b^2$. From these tables one can see that the transitions between states with the same partition are larger than transition between states with different $U(6)$ labels.

The isovector part $M1$ of the transition from 1^+ and higher energy 2^+ mixed symmetry states to ground state and first 2^+ state respectively is analyzed. The result is consistent with the identification of the 1^+ and the lowest 2_{ms}^+ in the these two nuclei. From the M1 operator, the isoscalar part is simply the angular momentum, M1 transition are determined by g_1 part only. For a good reproduction of calculated M1 transition we used the $g_1 = 2.8$ for the ^{48}Cr and ^{50}Cr . They lead to pure isovector M1 transitions. In ^{50}Cr isotope the calculated $B(M1; 1_1^+ \rightarrow 0_1^+)$ and $B(M1; 4_2^+ \rightarrow 4_1^+)$ values are in good agreement with the experimental data. We obtain $B(M1; 2_2^+ \rightarrow 2_1^+) = 0.6319 \mu_N^2$ which compares well with the $0.3938 \mu_N^2$ value observed experimentally. The $2_2^+ \rightarrow 2_1^+$ decay with its $E2/M1$ mixing ratio of $\delta = -0.03(6)$ experimentally is nearly pure M1 transition.

In ^{50}Cr the calculated 1^+ state at 8.901 MeV decays to ground state with reduced M1 transition $0.100 \mu_N^2$ and is in good agreement with experimental data of $0.0823(89) \mu_N^2$ reported in [41]. The relative B(E2) transitions between the states $J = 0^+, 1^+$ and 2^+ have different isospin values are calculated and summarized in Figs. 7 and 8 respectively.

V. CONCLUSION

Summarizing our results we may conclude that the IBM-3 description of the low-lying levels in the $^{48,50}\text{Cr}$ nuclei is satisfactory. We have shown, on the basis of energy levels and electromagnetic properties, the presence of mixed symmetry states near 5 and 3 MeV in $^{48,50}\text{Cr}$ respectively. In ^{48}Cr the 2_3^+ state at 5.069 MeV decays predominantly to the 2_1^+ state via a pure M1 transition, while in ^{50}Cr the 2_2^+ state decays through strongly M1 and weakly though E2 transition decay to 2_1^+ . These states are the candidate lowest mixed symmetry states in $^{48,50}\text{Cr}$ respectively. In our calculation the scissor mode 1^+ states in these nuclei lie higher than the lowest mixed symmetry 2^+ states, which were observed at 3.629 MeV in ^{50}Cr . No mixed symmetry state labeled with $[N-1, 1]$ and $T = 0$ exists in ^{48}Cr , and this is a natural result of group reduction. As a result, the scissor state in ^{48}Cr nucleus is higher than that in ^{50}Cr .

The calculated results are in good agreement with available experimental data, but more experiment data for those nuclei are needed to validate this nuclear structure predicted. In particular, the present calculations suggest a combined isospin and mixed symmetry excitation in the low-lying levels of these nuclei. The present calculation also predicts the existence of row-rowed partition state at about 9 MeV, and this type of states is typical of IBM3, not present in IBM1 and IBM2. It will be highly desirable to substantiate these predictions in future experiment.

Work is supported in part by National Natural Science Foundation of China under Grant No. 10047001, Major State Basic Research Development Program under Contract No. G200077400

-
- [1] A. Arima and F. Iachello, Ann. Phys. (N.Y) **99**(1976)253.
 - [2] A. Arima and F. Iachello, Ann. Phys. (N.Y) **111**(1978)201.
 - [3] A. Arima and F. Iachello, Ann. Phys. (N.Y) **123**(1979)468.
 - [4] F. Iachello and A. Arima, *The Interacting Boson Model*, Cambridge University Press, Cambridge, England (1987).
 - [5] W.D. Hamilton, A. Irbäck and J.P Elliott, Phys. Rev. Lett. **53** 2469(1984).
 - [6] B. Fazekas, T. Belgya, G. Molnar, A. Veres, R.A. Gatenby, S.W. Yates and T. Otsuka, Nucl. Phys. **A 548** (1992)249.
 - [7] I. Wiedenhover, A. Gelberg, T. Otsuka, N. Pietralla, J. Gableske, A. Dewald and P.von Brentano. Phys. Rev. **C56**(1997)R2354.

- [8] J.P. Elliott and A.P White, Phys. Lett **B 97**(1980)169.
- [9] G.L. Long Chinese Journal of Nuclear Physics, **16**(1994)331.
- [10] J. N. Ginocchio, Phys. Rev. Lett. **77** (1996) 28.
- [11] V.K.B. Kota, Phys. Lett **B 399**(1997)185.
- [12] V.S. Lac, J.P. Elliott and J.A. Evans, Phys. Lett. **B 394** (1997)231.
- [13] G. L. Long, C. Y. Gan and T. Y. Shen, Commun. Theor. Phys. **27** (1997) 317.
- [14] V.K.B. Kota, Ann. Phys. **265** (1998) 101.
- [15] P.O. Lipas, International Review of Nucl. Phys. Vol. **2** 1984, Edited by T. England, J. Rekestad and J-S Vadgen.
- [16] K. Langanke, P. Vogel, Dao-Chen Zhen, Nucl. Phys. **A 626** (1997)735.
- [17] S. Frauendorf and J. Sheikh, Nucl. Phys. **A 645** (1999)509.
- [18] Y. Durga Devi, Shadow Robmson, and Larry Zamick, Phys. Rev. **C 61** (2000)037305.
- [19] S.M. Fischer *etal.*, Phys. Rev. Lett.**87**(2002)132501.
- [20] P. Von Brentano, C. Frießner, R.N. Jolos, A.F.Lisetskiy, A.Schmidt, I.Schneider, N. Pietralla, T. Sebe and T. Otsuka Nucl. Phys.**A 704** (2002)115c.
- [21] E. Caurier ,J.L. Edido, G-Martinez-Pinedo, A.poves,J. Retamosa, L.M. Robledo and A.P. Zuker, Phys. Rev.Lett. **75**(1995)2466.
- [22] G. Martinez-Pinedo A.Poves, L.M. Robledo, E.Caurier, F.Nowacki, J.Retamosa and A. Zuker, Phys. Rev.**C 54**(1996)R2150.
- [23] J.A. Cameron G. ,J.L.Rodriguez,J. jonkman, G Hackman, S.M. Mullins, C.E. Svensson, J. C. Waddington, Lihong Yao, T.E. Drake, M.Cromaz, J. H. DeGraaf and G. Zwartz, H. R Andrews, G.ball, A. Galindo-Uribarri , V.P. Janzen, D. C. Radford and D. Ward, Phys. Rev.**C 58**(1998)808.
- [24] K. Langanke, Phys. Letter. **B 438**(1998)235.
- [25] Kenji Hara, Yang Sun and Takahiro Mizusaki, Phys. Rev. Lett. **83**(1999)1922.
- [26] S.M. Lenzi, D.R. Nagarajan, D. Bazzacco, D.M.Brink, M.A. Cardona, , G. de Angelis, M. de Poli, A.Gadea, D.Hojman, S. Lunardi, N.H.Medina, and C. Rossi Alvarez, Phys. Rev. **C 56**(1997)1313.
- [27] M. Hasegawa , K. Kaneko and S.Tazaki, Nucl. Phys. **A 674**(2000)411.
- [28] P.Descouvemont, Nucl. Phys. **A 709** (2002)275.
- [29] J.A. Evans, G.L. Long, J.P. Elliott, Nucl. Phys. **A 561**(1993)201.
- [30] J.A. Evans, J.P. Elliott, V.S. Lac and G.L. Long, Nucl. Phys. **A 593**(1995)85.
- [31] P.Van Isacker, unpublished.
- [32] G. Audi and A. H. Wapstra, Nucl. Phys. **A 565** (1993)1.
- [33] P.Van Isacker, K. Heyde, J. Jolie and A. Sevrin Ann.Phys.(N.Y) **171**(1986)253.
- [34] Hitoshi Nakada and Takaharu Otsuka, Phys. Rev. **C 55** (1997)2418.
- [35] S.A.A. Eid, W.D. Hamilton and J.P. Elliott, Phys. Lett. **B 166**(1986)267.
- [36] Hitoshi Nakada, Takaharu Otsuka and Takashi Sebe, Phys. Rev. Lett.**67**(1991)1086.
- [37] J.P. Elliott, J.A. Evans, V.S. Lac and G.L. Long, Nucl. Phys. **A 609**(1996)1.
- [38] J.P. Elliott, J.A. Evans and V.S. Lac, Nucl. Phys. **A 597** (1996)341.
- [39] V.S. Lac, unpublished (1996).
- [40] J. Lang, Krishna Kumar and J. H. Hamilton, Rev. of Mod. Phys., Vol.**54**, No. 1(1982) 119.
- [41] Richard B. Firestone, *Table of Isotopes*, Eighth Edition, edited by Virginia S. Shirley 1998.

TABLE I. The parameters of the IBM-3 Hamiltonian used for the description of the Cr-isotopes.

Nucleus	^{48}Cr	^{50}Cr
$\epsilon_{s\nu} = \epsilon_{s\pi}$	1.830	1.274
$\epsilon_{d\nu} = \epsilon_{d\pi}$	2.737	2.58
$A_i (i = 0, 1, 2)$	-6.264, -2.496, 2.496	-5.824, -2.108, 2.108
$C_{i0} (i = 0, 2, 4)$	-7.172, -5.018, -6.152	-5.802, -4.404, -6.042
$C_{i2} (i = 0, 2, 4)$	1.588, 3.742, 2.608	2.130, 3.528, 1.890
$C_{i1} (i = 1, 3)$	-2.993, -2.892	-2.370, -2.350
$B_i (i = 0, 2)$	-5.900, 2.860	-5.348, 2.584
$D_i (i = 0, 2)$	0.681, -0.814	0.890, -1.064
$G_i (i = 0, 2)$	-0.814, 2.496	-1.064, 0.890

TABLE II. Experimental [27,41] and calculated B(E2) (e^2B^2), calculated B(M1) (μ_N^2) and mixing ratio for ^{48}Cr isotope.

$J_i^+ \rightarrow J_f^+$	$B(E2)$		$B(M1)$		δ
	Exp.	Calc.	Calc.	Calc.	
$2_1^+ \rightarrow 0_1^+$	0.0320(41)	0.0309			
$2_2^+ \rightarrow 0_1^+$		0.0067			
$2_2^+ \rightarrow 2_1^+$		0.0007	0.0000		
$2_2^+ \rightarrow 0_2^+$		0.0101			
$1_1^+ \rightarrow 0_1^+$			0.6437		
$1_1^+ \rightarrow 0_2^+$			0.2362		
$1_1^+ \rightarrow 2_1^+$		0.0077	0.3923		-0.3141
$1_1^+ \rightarrow 2_2^+$		0.0091	0.2103		0.1434
$1_1^+ \rightarrow 2_3^+$		0.0016	0.1365		0.0517
$2_2^+ \rightarrow 4_1^+$		0.0166			
$4_1^+ \rightarrow 2_1^+$	0.0329(110)	0.0383			
$4_2^+ \rightarrow 2_1^+$		0.0012			
$6_1^+ \rightarrow 4_1^+$	0.0301(78)	0.0345			

 TABLE III. Experimental [27,40,41] and calculated B(E2) (in unit e^2b^2), B(M1) (in unit μ_N^2) and mixing ratio for ^{50}Cr isotope.

$J_i^+ \rightarrow J_f^+$	$B(E2)$		$B(M1)$		δ	
	Exp.	Calc.	Exp.	Calc.	Exp.	Calc.
$2_1^+ \rightarrow 0_1^+$	0.0217(25)	0.0247				
$2_2^+ \rightarrow 2_1^+$		0.0072	0.3938(716)	0.6319	$0.03^{+0.06}_{-0.04}$	0.1383
$2_2^+ \rightarrow 0_1^+$	0.0023(8)	0.0062				
$0_2^+ \rightarrow 2_2^+$		0.0385				
$2_2^+ \rightarrow 2_2^+$		0.0013		0.0238		-0.1291
$2_2^+ \rightarrow 0_1^+$		0.0109				
$2_3^+ \rightarrow 0_1^+$		0.0109				
$2_3^+ \rightarrow 2_1^+$		0.0209		0.0000		
$1_1^+ \rightarrow 0_1^+$			0.3300(400)	0.2778		
$1_1^+ \rightarrow 0_2^+$				0.0111		
$1_1^+ \rightarrow 2_1^+$		0.0072		0.2225		-0.2113
$1_1^+ \rightarrow 2_2^+$		0.0318		0.0013		1.2259
$1_1^+ \rightarrow 2_2^+$		0.0086		0.6559		0.0447
$1_2^+ \rightarrow 2_1^+$		0.0002		0.0026		0.2118
$1_2^+ \rightarrow 2_2^+$		0.0048		0.0012		4.0168
$3_1^+ \rightarrow 2_1^+$		0.1421		0.0221		0.4452
$3_1^+ \rightarrow 4_1^+$		0.0031		0.4741		0.0636
$4_1^+ \rightarrow 2_1^+$	0.0204(25)	0.0256				
$4_2^+ \rightarrow 4_1^+$		0.0110	0.1325(375)	0.4885	$-0.02^{+0.16}_{-0.52}$	0.1281
$5_1^+ \rightarrow 3_1^+$		0.0031				
$5_1^+ \rightarrow 4_1^+$	0.0001	0.0171		0.0033	-0.47(16)	0.5005
$6_1^+ \rightarrow 4_1^+$	0.0235(47)	0.0165				




FIG. 1. Comparison between lowest excitation energy bands ($T = T_z, T_z + 1$ and $T_z + 2$) of the IBM-3 calculation and experimental excitation energies of ^{48}Cr . The experimental data are taken from Ref. [41].




FIG. 2. Comparison between lowest excitation energy bands ($T = T_z, T_z + 1$ and $T_z + 2$) of the IBM-3 calculation and experimental excitation energies of ^{50}Cr . The experimental data are taken from Ref [41].




FIG. 3. The variation in level energy of ^{48}Cr as a function of C_{11} ; all the other parameters were kept at their best-fit values.




FIG. 4. The variation in level energy of ^{48}Cr as a function of C_{31} ; all the other parameters were kept at their best-fit values.




FIG. 5. The variation in level energy of ^{50}Cr as a function of C_{11} ; all the other parameters were kept at their best-fit values.




FIG. 6. The variation in level energy of ^{50}Cr as a function of C_{31} ; all the other parameters were kept at their best-fit values.




FIG. 7. Energy levels and relative B(E2) for ^{48}Cr isotope.




FIG. 8. Energy levels and relative B(E2) for ^{50}Cr isotope.